
# **Plants of South Eastern New South Wales**



Flow ering stems. Photographer Don Wood, Nadgee State Forest south of Eden.



Flow ering stems. Australian Flant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra



Line drawing. b. flowering branch. Inset: detail of veins in 'leaf'. MMbir, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

## Common name

Bower wattle, River wattle, Narrow-leaf bower wattle

### **Family**

Fabaceae

#### Where found

Dry forest, woodland, and moist gullies. Coast and ranges south of Wollongong. Planted, and an occasional garden escape, in Canberra and Queanbeyan.

#### **Notes**

Tree or shrub to 10 m tall. Fleshy seed stalks/arils. Bark smooth. Branchlets ridged, hairless to slightly hairy, often sticky, arching to weeping. 'Leaves' alternating up the stems, usually weeping, 4-10 cm long, 1-4 mm wide, sparsely hairy to hairless with fringed margins, usually with 3 main longitudinal veins (midvein most prominent), dotted with resin glands, often sticky, tips pointed with a mucro. Flower heads pale yellow, globular, 10-25 flowered (easiest seen in late buds), 3-6 mm in diameter, in clusters of 1-2 flower heads. Flowers Winter to Spring.

Intermediate forms between narrow-leaved Acacia cognata and broader-leaved Acacia subporosa occur.

Family was Mimosaceae.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Acacia~cognata (accessed 29 April 2021)

World Wide Wattle line drawings, photos, and description: <a href="http://www.worldwidewattle.com/imagegallery/image.php?">http://www.worldwidewattle.com/imagegallery/image.php?</a>
<a href="p=0&l=c&id=8928&o=1">p=0&l=c&id=8928&o=1</a>

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)