Plants of South Eastern New South Wales




Flow ering stem Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra



Flowering stem Photographer Don Wood, Quumbean Nature Reserve near Queanbeyan



Shrub. Australian Plant Image Index, photographer Colin Totterdell, Jerrabombera NSW



Line drawing. c. flowering branch. Inset: detail of veins in 'leaf'. MMbir, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Poverty wattle, Mitta wattle, Dawson's wattle

Family

Fabaceae

Where found

Dry forest, woodland, and grassy areas. Western Slopes, ACT, and tablelands.

Notes

Shrub or tree to 4 m tall. Branchlets with prominent resinous ribs, hairy with minute hairs. 'Leaves' alternating up the stems, 3.5-12 cm long, 2-8 mm wide, hairless, more or less resinous, with up to about 10 longitudinal veins (1 or 2 being more prominent), tips somewhat pointed with a mucro. 1 marginal gland near the base. Flower heads yellow, globular, 4-10 flowered (easiest seen in late buds), 2-5 mm in diameter, in elongated clusters of 2-9 flower heads. Flowers Winter to early Summer. Pods linear, shiny.

Family was Mimosaceae.

Regarded as indigenous and naturalised in the ACT. All native plants on unleased land in the ACT are protected.

Vulnerable Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Acacia~dawsonii (accessed 29 April 2021)

World Wide Wattle line drawings, photos, and description: http://www.worldwidewattle.com/imagegallery/image.php?
p=0&l=d&id=23624&o=1

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)