Plants of South Eastern New South Wales




Flow ering stem Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra



Pods and 'leaves'. Photographer Jackie Mles



Open pods and 'leaves'. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra



Line drawing. g. leafy branch; pod. Inset: detail of 'leaf'. MMbir, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Woolly bear wattle, Lucas's wattle

Family

Fabaceae

Where found

Dry forest, woodland, and heath, often in rocky areas. Tablelands and ranges south from north east of Cooma.

Notes

Shrub to 4 m tall. Fleshy seed stalks/arils. Branchlets more or less cylindrical, hairy. 'Leaves' alternating up the stems, 1.5-6 cm long, 6-15 mm wide, straight or slightly curved, surfaces hairy, midvein and marginal veins prominent, margins flat to slightly wavy, tips blunt with a mucro, one side of the base sometimes longer than the other. Flower heads yellow, oval to globular, 5-10 mm long, 12-40 flowered (easiest seen in late buds), in elongated clusters of 1–10 flower heads, the axis 1–5 cm long. Individual flower stalks 1–10 mm long. Flowers Winter to Summer. Pods rusty brown velvety.

Family was Mimosaceae.

Vulnerable Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Acacia~lucasii (accessed 27 April 2021)

World Wide Wattle photos, line drawings and description: http://www.worldwidewattle.com/imagegallery/image.php?
p=0&l=l&id=23726&o=1

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)