Plants of South Eastern New South Wales



Rowers. Australian Rant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Flowering branch. Australian Flant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Seed cases and leaves. Photographer Jackie Mles, Byron Bay



Shrub. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Trunk. Photographer GLemann, Bawley Point

Common name

Brush kurrajong

Family

Malvaceae

Where found

Rainforest, wet forest, clearings, and roadsides. Coast and ranges north of the Bermagui district.

Notes

Small tree or shrub to 8 m high. Bark brown to greyish, smooth with horizontally elongated lenticels. Stems hairy with pale or brown stellate hairs about 0.5 mm long (needs a hand lens or a macro app on your phone/tablet to see), lenticels not obvious at first. Leaves alternating up the stems, 5-20 cm long, 20-120 mm wide, lower surface covered with white or or pale brown centred stellate hairs, margins irregularly toothed, sometimes with short lobes. Flowers cream to white, sometimes pinkish, 9-11 mm in diameter, with 5 petals and 5 staminodes that look like petals, as long as the petals. Flowers in branched clusters 3.5-5 cm long.

Family was Sterculiaceae.

Difficult to distinguish from *Androcalva rossii* in southern NSW. It has smaller leaves, and smaller stellate hars on the branchlets, leaves, and flower clusters.

Author: Betty Wood.

 $This identification \ key \ and \ fact \ sheets \ are \ available \ as \ a \ free \ mobile \ application:$





Creative Commons Attribution 3.0 Australia (CC BY)