
Plants of South Eastern New South Wales



Seeding plants. Australian Plant Image Index, photographer Colin Totterdell, Kosciuszko National Park



Line drawings (as *Euchiton nitidulus*). f. plant; flower head. EMayfield, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board.

Common name

Shining cudweed

Family

Asteraceaae

Where found

Heath, grassland, herbfields, and near streams and bogs. Kosciuszko National Park. One record on the tablelands from between Jindabyne and Dalgety.

Notes

Mat- or cushion-forming **perennial** herb to 0.03 m high, rooting from the **nodes**. Flower stalks **woolly**, densely covered at the base with the dried remains of old leaves. Leaves crowded toward the branch-tips, 0.5–0.7 cm long, 2–3 mm wide, covered with shining, silvery to rust-coloured **appressed** hairs on both surfaces. Flower **heads** about 9-10 mm in diameter, with 0 **petals**. **Bracts** surrounding the flower heads brownish, with long hairs at the base, hairless, shining, dry and translucent in the upper half. Inner btsscts 7-9 mm long. Flower heads single. Flowering: December –March

Vulnerable Australia. Vulnerable NSW. Provisions of the Biodiversity Conservation Act 2016 No 63 relating to the protection of protected plants generally also apply to plants that are a threatened species.

Rare Vic.

NSW Threatened Species profile: http://www.environment.nsw.gov.au/threatenedspeciesapp/profile.aspx?id=10320 (accessed 3 January, 2021)

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Argyrotegium~nitidulum (accessed 3 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:







Creative Commons Attribution 3.0 Australia (CC BY)