Plants of South Eastern New South Wales



Flowering stems. Australian Flant Image Index, photographer Murray Fagg, near Cow ra



Stem, leaves, and seed cases. Australian Flant Image Index, photographer Murray Fagg, near Cowra



Flowering plant. Australian Flant Image Index, photographer Murray Fagg, near Yass

Common name

Tarvine, Murra, Tah-vine

Family

Nyctaginaceae

Where found

Wide range of dry habitats. Mainly Western Slopes. Occasionally elsewhere.

Notes

Perennial herb, prostrate to erect, stems to 1m long or more. Stems hairless or hairy to sticky hairy. Leaves opposite each other, 1-4 cm long, 3-20 mm wide, hairy, sometimes sticky, sometimes sparsely warty on the upper surface, margins flat or sometimes slightly wavy, and entire or with small teeth, tips pointed, blunt, or notched. Flowers about 3 mm wide, pink, white or mauve, bell shaped, with 5 lobes, in clusters of 2–6. Flowers Summer to Autumn.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Boerhavia~dominii (accessed 4 May, 2021)

Author: Betty Wood.

 $This \, identification \, key \, and \, fact \, sheets \, are \, available \, as \, a \, \, free \, mobile \, application: \, \\$







Creative Commons Attribution 3.0 Australia (CC BY)