Plants of South Eastern New South Wales



Flowering stems. Photographer Don Wood, Mungerarie State Forest west of Moruya



Open seed cases. Australian Flant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Shrub. Australian Plant Image Index, photographer S Donaldson, Caves Road, NSW

Common name

Showy Boronia, Sydney Boronia, Ledum Boronia

Family

Rutaceae

Where found

Dry forest, shrubland, heath, and near streams. Coast, ranges, and the eastern edge of the tablelands.

Notes

Shrub to 2.5 m high. Branchlets stellate-hairy (needs a hand lens or a macro app on your phone/tablet to see), often glandular-warty. Leaves aromatic when rubbed, opposite each other, simple, or compound with 3–7 leaflets, terminal leaflet usually the longest. Simple leaves 0.3–4.3 cm long, 1–7 mm wide. Compound leaves with a winged rachis 0.2–0.8 cm long, leaflets 0.4–3.4 cm long, 1–5 mm wide. Leaves/leaflets with the upper surface sparsely hairy to hairless and shining, lower surface hairy and paler, margins entire and curved to rolled down, tips pointed to blunt. Flowers pale to bright pink, occasionally white, densely hairy, with 4 petals each 5–12 mm long, persistent at the seeding stage. Petal margins touching each other in bud. 8 stamens. Flowers in 1-3 flowered clusters. Flowers winter-summer.

Protected NSW.

Vulnerable Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Boronia~ledifolia (accessed 5 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)