## **Plants of South Eastern New South Wales**



Flow ering stems. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT

Flow ering stems. Photographer Don Wood, Nadgee Nature Reserve



Shrub. Australian Flant Image Index, photographer R Hotchkiss, Australian National Botanic Gardens, Canberra, ACT

## **Common name**

Forest Boronia

Family

Rutaceae

## Where found

Forest, woodland, and heath. Coast and ranges south of Eden.

## Notes

Tree or shrub to 7 m high. Branchlets hairless, or sparsely hairy in 2 lines separated by hairless leaf bases continuing down the stem, and glandular-warty. Leaves aromatic when rubbed, opposite each other, 4.5–7 cm long, 20–42 mm wide, compound, with 7–17 leaflets, the terminal leaflet shortest. Rachis 1.3–3.5 cm long, leaflets 0.5–2.5 cm long, 1–4.5 mm wide, upper surface hairless and more or less glandular-warty, lower surface paler than the upper surface, margins finely toothed and curved down, tips pointed. Flowers deep to pale pink, often brighter pink on the outside, occasionally white, with 4 petals each 5–9 mm long. Petals overlapping each other in bud, falling early. 8 stamens. Flowers in 1–15-flowered clusters.

Protected NSW.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Boronia~muelleri (accessed 5 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:







iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)