Plants of South Eastern New South Wales



Flow ering stems. Australian Flant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Flow ering stems. Photographer Don Wood, Eurobodalla Regional Botanic Gardens, north of Moruya

Common name

A boronia

Family

Rutaceae

Where found

Forest in damp shady situations. North from Moruya. Coast, ranges, and eastern edge of the tablelands.

Notes

Tree or shrub to 4 m high. Branchlets with 2 grooves separated by leaf bases continuing down the stems, and hairless. Leaves aromatic when rubbed, opposite each other, compound, with 3–15 leaflets, the terminal leaflet often the shortest. Rachis 0.9–4 cm long, slightly winged, leaflets 0.5–3 cm long, 1–6 mm wide, lower surface slightly paler than the upper surface, margins finely glandular-scalloped and rolled to curved down, tips pointed. Flowers pink, with 4 petals each 6–9 mm long. Petals overlapping each other in bud, falling early. 8 stamens. Flowers in 2–6 flowered clusters. Flowering: August–November.

Protected NSW.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Boronia~thujona (accessed 5 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:







Creative Commons Attribution 3.0 Australia (CC BY)