


Plants of South Eastern New South Wales



Flowering plant. Australian Plant Image Index, photographer Alexander Schmidt-Lebuhn, Koorawatha Nature Reserve south of Cowra



Flowering plant. Australian Plant Image Index, photographer Alexander Schmidt-Lebuhn, Illunie Nature Reserve south of Cowra



Line drawings. b. flowering branch; seed (front and side view). Gloria Thonlinson, Daisy Study Group, © 2021 Royal Botanic Gardens Board, Melbourne, Vic

Common name

Dookie Daisy

Family

Asteraceae

Where found

Woodland. Rocky areas and moist sites. Western Slopes.

Notes

Annual herb to 0.35 m high or sprawling, hairy to slightly hairy and somewhat sticky. Leaves basal and **alternating** up the stems, basal leaves usually not persisting. Leaves 0.4–9 cm long, 0.5–2.5 mm wide, deeply lobed to **dissected**, stem-clasping, tips blunt. Flower **heads** with 8–12 white '**petals**' about 3.5–5 mm long, and yellow centres to 7 mm in diameter. 8–14 **bracts appressed** to the flower heads. Flower heads usually single. Flowering: September–November.

Definite identification of most species of *Brachyscome* depends on microscopic examination of the seeds.

Vulnerable Vic. Listed in the Flora and Fauna Guarantee Act, Vic.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Brachyscome~gracilis> (accessed 6 January, 2021)

Author: Betty Wood.



Creative Commons Attribution 3.0 Australia (CC BY)

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition