
Plants of South Eastern New South Wales



Row ering stem Australian Rant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Line drawings. a. upper stem leaf; b. seed case. M Mbir, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Cauliflower, Collards

Family

Brassicaceae

Where found

Occasionally naturalised. Mainly Sydney area. Occasionally elsewhere.

Notes

Introduced biennial or perennial herb to 3 m high. Leaves more or less fleshy. Stems hairless, becoming woody, with conspicuous basal leaf scars. Leaves basal and alternating up the stems, glaucous, hairless. Lower leaves to 40 cm long, stalked, broad and rounded, margins variously scalloped, occasionally lobed; upper leaves not lobed, stalkless, stem-clasping, bract -like. Flowers yellow, with 4 petals 10–25 mm long. Flowers in elongated clusters. Flowers Sep.–Nov. Seed cases spreading horizontally, then becoming erect, 5–10 cm long, including the beak, 2–5 mm wide, cylindrical, or weakly compressed. Beak 5–11 mm long, with 1 or 2 seeds. Stalk 10–30 mm long. Seeds dark brown to black.

Fruiting plants. Photographer Greinsteidl Germany

Cultivated as a vegetable.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Brassica~oleracea (accessed 6 January, 2021)

Author: Betty Wood.



This identification key and fact sheets are available as a free mobile application:



Android edition iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)