Plants of South Eastern New South Wales



Rowering plant. Photographer Jackie Mles, Kosciuszko National Park



Rowering plants. Australian Rant Image Index, photographer Dave Mallinson, Boboyan, ACT



Flow ering plant. Photographer Don Wood, Kow en Stock Route Reserve, ACT



Line drawings. g. leaf; seed. EMayfield, National Herbariumof Victoria, ©2021 Royal Botanic Gardens Board

Common name

Rough burr-daisy, Blue burr-daisy

Family

Asteraceae

Where found

Forest, woodland, and grassland. Kosciuszko National Park, the mountains to the north, ACT, tablelands, and the western edge of the ranges. Occasionally on the Western Slopes.

Notes

Perennial herb to 0.45 m high, rhizomatous or stoloniferous, forming colonies. Stems sparsely hairy to hairy or bristly. 'Seeds' flattened, 2.5–4 mm long, smooth, hairless, with prominent marginal ridges and 4-6 spines 1–4 mm long, barbed at the tips, often with smaller spines 1 mm long, all hairy at the base. 'Seeds' stick to clothing. Basal leaves 3–20 cm long, 7–40 mm wide, broadly spoon-shaped, margins toothed or dissected. Stem leaves alternating up the stems, 0.5–3 cm long, 2–10 mm wide. Leaves rough or hairy, bases slightly stem-clasping. Flower heads 15–35 mm in diameter, with 10-65 white, mauve, magenta, or purplish 'petals' 7–13 mm long, and yellow centres. Flowering: throughout the year.

Definite identification of all species of Calotis is made by examining the burrs.

All native plants on unleased land in the ACT are protected.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Calotis~scabiosifolia (accessed 7 January, 2021)

Author: Betty Wood.

6 ۲ This identification key and fact sheets are available as a free mobile application:



Android edition



Creative Commons Attribution 3.0 Australia (CC BY)