

Plants of South Eastern New South Wales



Flower and leaves. Photographer Richard Hartland, New Zealand



Flower and leaves. Photographer Richard Hartland, New Zealand



Flowering stems. Photographer Don Wood, Barragoot Lake south of Barragui

Common name

Sea Bindweed

Family

Convolvulaceae

Where found

Coastal dunes and rocky areas.

Notes

Perennial herb with trailing or rarely twining stems. Hairless. Leaves somewhat fleshy, **alternating** along the stems, 1.5–4 cm long, 20–60 mm wide, more or less kidney shaped, margins wavy. Flowers white, pink or purplish, 30–50 mm long, trumpet-shaped, with 5 lobes. **Stigma** with 2 lobes. Sepals free from each other. **Bracteoles** broadly oval, tips blunt to rounded, 10–15 mm long, slightly shorter than the **sepals**. Flowers single. Flowers Spring to Summer.

Vulnerable Vic.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lv=sp&name=Calystegia~soldanella> (accessed 7 January, 2021)

Author: Betty Wood.



Creative Commons Attribution 3.0 Australia (CC BY)

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition