


Plants of South Eastern New South Wales



Line drawings. m. basal leaves. MIMbir, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Franklin Bitter-cress

Family

Brassicaceae

Where found

Subalpine areas. ACT, the mountains to the west, Kosciuszko National Park, and tablelands south of the ACT.

Notes

Erect perennial herb to 0.3 m high, with a taproot. Basal leaves to 15 cm long, **entire** and spoon-shaped or sometimes **compound**, forming a more or less dense **rosette**. Stem leaves 0-3, **alternating** up the stems, 1-5 cm long, compound, becoming smaller in size up the stem. Flowers with 4 **petals** each 4.5-7 mm long, all white, pink, or white inside and pink on the outside. 6 **stamens**. Flowers usually in many-flowered clusters. Flowers spring.

All native plants on unleased land in the ACT are protected.

Endangered Vic. Listed in the Flora and Fauna Guarantee Act, Vic.

PlantNET description with line drawings and photos of dried specimens: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Cardamine~franklinensis> (accessed 7 January, 2021)

Author: Betty Wood.



Creative Commons Attribution 3.0 Australia (CC BY)

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition