
# **Plants of South Eastern New South Wales**



Flowers and leaf. Photographer Chris Clarke, Anakie, Vic



Plant. Photographer Chris Clarke, Anakie, Vic



Line drawing. g. plant. MMbir, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

#### Common name

Forest bitter-cress

## **Family**

Brassicaceae

### Where found

Hilly to mountainous forested areas. Mainly Kosciuszko National Park, the mountains to the north, and ACT. Occasional elsewhere.

#### **Notes**

Small annual or perennial herb to 0.25 m high. Stems hairless or covered with minute warts. Basal leaves 2-7 cm long, simple, or compound with 3-5 leaflets, forming a persistent rosette. Lateral leaflets more or less round, terminal leaflet oval to cordate. Stem leaves, 0-3, alternating up the stems, 0.5–6 cm long, variable in shape, upper leaves much reduced, margins sometimes warty. Flowers with 4 petals each 3–5 mm long, white, pink, or white inside, pink outside. 6 stamens. Flowers usually in few-flowered clusters. Flowers Spring-Autumn.

All native plants on unleased land in the ACT are protected.

Vulnerable Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Cardamine~papillata (accessed 7 January, 2021)

Author: Betty Wood.

 $This identification \ key \ and \ fact \ sheets \ are \ available \ as \ a \ free \ mobile \ application:$ 





