Plants of South Eastern New South Wales



Flow ering stem. Australian Flant Image Index, photographer Murray Fagg, Shepherds Lookout, ACT



Rowering plant. Australian Rant Image Index, photographers RG & FJ Richardson, near Warrnamboool, Vic



Rosette. Australian Plant Image Index, photographer JM DiTomaso, California, USA

Common name

Slender thistle

Family

Asteraceae

Where found

Dry forest, woodland, shrubland, pastures, roadsides, wasteland and cultivation, and rocky sites. Widespread.

Notes

Introduced annual herb to 2 m high, initially forming a basal rosette. Stems grooved, initially cobwebby, sometimes almost hairless, with spiny wings. Wings more or less discontinuous, and broken into spiny lobes to 5 mm wide. Leaves with spiny margins. Flower heads surrounded by spiny bracts. Basal leaves 5–25 cm long, 20–80 mm wide, upper surface green and hairy, lower surface white and cobwebby, deeply dissected. Stem leaves similar, alternating up the stems, 5–15 cm long, 20–70 mm wide. Flower heads cylindrical, to about 20 mm long. Florets 10-17 mm long, purple-pink. Flower heads in clusters of 1-4. Flowering: Oct.—Nov.

Pest plant ACT. Noxious weed Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Carduus~pycnocephalus (accessed 7 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





