


Plants of South Eastern New South Wales



Flowering stem. Photographer Don Wood, Moruya State Forest near Moruya



Shrub. Australian Plant Image Index, photographer Murray Fagg, between Wauchope and Walcha



Line drawings. f. flowering branch; flower head; leaf. E Beckett, E. University of Oxford, © 2021 Royal Botanic Gardens Board, Melbourne, Vic

Common name

Three-veined cassinia

Family

Asteraceae

Where found

Forest, shrubland, roadsides and rocky sites. Coast, ranges, and the eastern edge of the tablelands.

Notes

Shrub or tree to 8 m high. Stems yellowish, greyish, or whitish, resinous, sticky hairy, with fine **gland**-tipped hairs. Older stems more sparsely hairy. Leaves **alternating** up the stems, 3–12 cm long, 3–8 mm wide, upper surface hairless or slightly sticky, lower surface sticky hairy and with 3 prominent more or less parallel **veins**, especially at the base, tips pointed, margins flat or curved down. Flower **heads** white, oblong to bell shaped at maturity, 2.5–3.5 mm long, 1–1.5 mm wide, with 3 or 4 creamy-white **florets**, in a dense cluster of 300–500 flower heads 60–120 mm in diameter. Flowering: summer–autumn.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Cassinia-trinerva> (accessed 7 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)