# **Plants of South Eastern New South Wales**



Juvenile plant. Photographer Peter Woodard, Sydney



Base of trunk. Australian Plant Image Index, photographer Murray Fagg, TooloomNational Park, NSW, east of Warwick, Qld

## Common name

Churnwood, Silky beech, Soapbox

# **Family**

Icacinaceae

### Where found

Rainforest and moist sheltered gullies. Coast and ranges north of Batemans Bay.

#### **Notes**

Tree to 50 m high. Fruit fleshy. Trunk channelled. Bark light to grey-brown, finely fissured and corky. Branchlets with appressed brownish hairs, soon becoming hairless. Leaves alternating up the stems, 5–15 cm long, 30–70 mm wide, hairless, upper surface glossy, lower surface paler, margins entire, domatia on the lower surface small, pocket-like, in the vein angles along the midrib (handlens/macro app on your mobile phone needed). Male and female flowers on different plants. Flowers white, or creamy green to yellow, with 5 petals each 4–5 mm long, Flower clusters 5–15 cm long. Flowering: May–September. Fruit black, globular to oval, 15-25 mm long.

Family Cardiopteridaceae in PlantNET.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Citronella~moorei (accessed 19 April 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:







Creative Commons Attribution 3.0 Australia (CC BY)