# **Plants of South Eastern New South Wales**



Flow ering branch. Photographer Don Wood, Booderee Botanic Gardens, Jervis Bay



Tree. Photographer Raffi Kojian, Royal Botanic Gardens Sydney



Trunk. Photographer Peter Woodard, Wyrrbalong National Park near The Entrance

## Common name

Corkwood, Eye opening tree, Poisonous corkwood, Poison corkwood, Yellow basswood, Duboisia

### **Family**

Solanaceae

#### Where found

Forest, rainforest regrowth, and wet gullies. Coast and ranges, north of Batemans Bay.

#### **Notes**

Tree or shrub to 24 m high. Fruit fleshy. Bark creamy brown, soft, thick and corky, strongly fissured on older plants. Young stems densely hairy with stellate hairs (needs a hand lens or a macro app on your phone/tablet to see), becoming hairless. Leaves alternating up the stems, 4–15 cm long, 10–40 mm wide, very soft, hairless, dull. Flowers white with purple striations in the throat, 4–8 mm long, tubular, the tube bell-shaped, 2–4 mm in diameter at the mouth, with 5 lobes each 1–3.5 mm long. Flowers in open many-flowered branched clusters. Flowering: mostly winter to spring. Fruit purple to black, usually globular, 4.5-8 mm in diameter. All parts of the plant poisonous.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Duboisia~myoporoides (accessed 12 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





