
Plants of South Eastern New South Wales




Rowers and leaves. Australian Rant Image Index, photographer Murray Fagg, Cooma NSW



Pant. Photographer Don Wood, near Namadgi Visitors Centre near Tharwa, ACT



Pant. Australian Pant Image Index, photographer Murray Fagg, Scottsdale Bush Heritage Reserve, near Bredbo



Line drawings (as *Chenopodium pumilio*). m leaf; seed case. MMbir, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Small crumbweed

Family

Chenopodiaceae

Where found

Disturbed sites. Widespread.

Notes

Annual or short lived perennial herb to 0.25 m tall or sprawling to prostrate. Stems hairy, slightly sticky. Leaves aromatic when rubbed, alternating along the stems, 0.5-2 cm long, 3–10 mm wide, hairy, sticky, margins entire, toothed, lobed, or wavy. Flowers female or bisexual, tiny (less than 3 mm in diameter) with 5 'petals' united towards the base, in small clusters. Seed cases whitish when mature.

Was Chenopodium pumilio.

All native plants on unleased land in the ACT are protected.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Dysphania~pumilio (accessed 12 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:

é

iOS edition

Android edition



Creative Commons Attribution 3.0 Australia (CC BY)