Plants of South Eastern New South Wales



Flow ering stems. Photographer Don Wood, Coolangubra State Forest near Bombala



Tree. Australian Plant Image Index, photographer Murray Fagg, near Bemboka



Fruiting stems. Photographer Don Wood, SE Forests National Park



Line drawings. EMMayfield, a. flowering branch; b. flower; c. fruit. National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Black Olive Berry, Mountain Blueberry, Mountain Quandong

Family

Elaeocarpaceae

Where found

Rainforest and wet gullies. Coast, ranges and tablelands.

Notes

Tree or **shrub** to 25 m tall. Fruit fleshy. Trunk often flanged towards the base. Bark dark grey, relatively smooth with some fissures and wrinkles, and with horizontally elongated **lenticels**. Young stems densely hairy with rusty to pale hairs, becoming hairless. Leaves **alternating** up the stems, 3–11 cm long, 10–30 mm wide, tough to very stiff, upper surface hairless, lower surface hairy with no **domatia**, margins distinctly toothed, no joint at the top of the leaf stalk. Flowers white to cream, sometimes flushed with pink, with 4-5 **petals** each 6-12 mm long, **entire** or shallowly lobed, not fringed. Flower clusters about 1.5-6 cm long. Flowers Spring to Summer. Fruit at first with a waxy bloom, ripening purple or blue-black to glossy black, oval, about 6 mm in diameter. Fruits Summer to Autumn.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Elaeocarpus~holopetalus (accessed 12 January, 2021)

Author: Betty Wood.



This identification key and fact sheets are available as a free mobile application:



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)

(cc