Plants of South Eastern New South Wales



Line drawings. d. plant; seed; seeding flower head. M Moir, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Daisy Fleabane

Family

Asteraceae

Where found

Woodland and rocky sites. Kosciuszko National Park.

Notes

Perennial herb to about 1 m high, with annual stems. Stems hairless, minutely sticky hairy below the flower heads. Leaves alternating up the stems, 2–13 cm long, 3–12 mm wide, margins sometimes purplish, margins entire or the lowermost leaves toothed. Flower heads with many pale purple to white 'petals' in one to several rows, each 1.5–8 mm long, more or less erect. Flower heads below the 'petals' 3-8 mm long, 10-20 mm in diameter, hemispherical or bell-shaped. Flower heads in open clusters, appearing single. Flowering: Jan.–Apr.

Vulnerable Vic.

PlantNET description with line drawings: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Erigeron~conyzoides (accessed 14 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:







Creative Commons Attribution 3.0 Australia (CC BY)