


Plants of South Eastern New South Wales



Flower head and seeding flower head. Photographer Jackie Miles



Flowering plants. Photographers Forest & Kim Starr, Hawaii



Line drawings. c. plant; flower head at left. M.Mbir, National Herbarium of Victoria, ©2021 Royal Botanic Gardens Board

Common name

Bony-tip Fleabane, Seaside Daisy

Family

Asteraceae

Where found

Garden escape from gardens and rockeries. Moist disturbed sites and along streams. Coast, ranges, and the eastern edge of the tablelands, north of Nowra. Canberra. Occasionally elsewhere.

Notes

Introduced sprawling **perennial** herb to 0.4 m high, much-branched from the base. **Rhizomatous**. Stems more or less **wiry**. Leaves **alternating** up the stems. Lower leaves 1–6 cm long, 15–30 mm wide, 3–7 lobed or coarsely 3–5-toothed at the tips; grading to the upper leaves 0.5–2.5 cm long, 1–5 mm wide, margins **entire**, often with a hardened tip. Flower **heads** with many **petals** in several rows, pale purple, turning white and finally pink or red; each 3–10 cm long, and yellow centres 8–15 mm in diameter with many **florets**. Flower heads below the '**petals**' approximately hemispherical. Flowering: Oct.–Feb.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Erigeron~karvinskianus> (accessed 25 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)