Plants of South Eastern New South Wales



Flowering and seeding stems. Photographer Don Wood, Flonacle Reserve, Canberra, ACT



Flowering plant. Photographer Don Wood, Scotia Sanctuary, western NSW



Plant. Photographer Don Wood, Scotia Sanctuary, western NSW

Common name

Blue storksbill, Native crowfoot, Blue crowfoot, Blue heron's-bill

Family

Geraniaceae

Where found

Dry forest, woodland, shrubland, grassy areas, and disturbed sites. Widespread. Rare on coast and ranges south of the Sydney area..

Notes

Annual or biennial to perennial herb, to 0.9 m tall or sprawling. Stems with scattered long stifff hairs. Leaves basal and opposite each other, basal leaves 1.5-6 cm long, 10-30 mm wide, stem leaves smaller. Leaves hairy, deeply dissected into three flat lobes which are again lobed or toothed to scalloped. Leaves sometimes appear compound, with 3 leaflets. Flowers blue, with 5 petals, each about 6-10 mm long, in clusters of 2-6 flowers, rarely single. Flowers most of the year.

All native plants on unleased land in the ACT are protected.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Erodium~crinitum (accessed 14 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





