Plants of South Eastern New South Wales




Flower, buds, and leaves. Australian Flant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra ACT



Burr. Photographer Russell Best, Kosciuszko National Park



Flow ering plants. Photographer Don Wood, Namadgi National Park, ACT



Line drawings. I. leaf; m flower; n. burr. BMMayfield, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Herb Bennet, Common Avens, Avens

Family

Rosaceae

Where found

Along streams and on the margins of and in bogs in montane to alpine areas. ACT, the mountains to the west, Kosciuszko National Park, and nearby. Occasionally on the western edge of the ranges/eastern edge of the tablelands.

Notes

Perennial herb to 1 m high, with a short rhizome. Stems reddish green, hairy to hairless. 'Seeds' forming burrs with many reddish hooked spines that stick to clothing. Leaves basal and alternating up the stems, 4–30 cm long, decreasing in size up the stems, hairy on both surfaces, mainly along the veins, compound with 3–11 leaflets, or dissected, the segments irregularly and deeply toothed. Stipules leaflike, toothed and mainly 3-lobed, 0.75–2 cm long. Flowers yellow, with 5 petals each 4.5–8 mm long. Flowers in few-flowered open clusters, appearing single. Flowering: summer.

Varieties not recognised in NSW.

All native plants on unleased land in the ACT are protected.

PlantNET description (as *Geum urbanum*): http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Geum~urbanum (accessed 13 April 2021) Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)