

Plants of South Eastern New South Wales



Flower. Photographer Jackie Mles, Currowan




Flowering plant. Photographer Don Wood, Wadbilliga National Park



Stems with flower and seeds. Photographer Jackie Mles, Bungonia



Seeds. Photographer Andre Messina, unknown place, ©2021 Royal Botanic Gardens Board, Melbourne



Line drawing. In VICFLORA

Common name

Cobbler's tack

Family

Asteraceae

Where found

Wide range of **habitats**. Coast, ranges, and the eastern edge of the tablelands, north from Wadbilliga National Park.

Notes

Perennial herb, stems woody at the base, to 0.60 m tall or sprawling. 'Seeds' surmounted by two stiff barbed **awns**. Almost hairless. Leaves mainly in a basal **rosette**, a few **alternating** up the stems. Basal leaves 1–12 cm long, 5–50 mm wide, green above, **glaucous** below, hairless, sometimes undivided and **linear**, usually deeply divided into narrow **segments**. Stem leaves becoming smaller and linear higher up the stem. Flower **heads** 5–8 mm in diameter overall, with 5 to about 20 toothed yellow '**petals**', or with 0 '**petals**', and yellow centres with 7–12 **florets**. Flower heads single. Flowers Spring to Autumn.

Was *Glossogyne tannensis*.

Vulnerable Vic.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lv=sp&name=Glossocardia-bidens>
(accessed 18 January, 2021)

Author: Betty Wood.



Creative Commons Attribution 3.0 Australia (CC BY)

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition