Plants of South Eastern New South Wales



Flower. Photographer Jackie Miles



Flowering stems. Photographer Jackie Mles



Line drawing. g. flowering branch. EMayfield, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Lanky Goodenia

Family

Goodeniaceae

Where found

Forest, often in damper sites. Widespread south of the Hume and Sturt Highways.

Notes

Short-lived perennial herb to 0.50 m high. Stems with spreading or appressed simple hairs, or hairless. Leaves in a loose basal rosette, mostly alternating up the stems, 2–9 cm long, 4–20 mm wide, tips pointed to blunt, surfaces hairy to becoming hairless, margins toothed to entire. Flowers 12–20 mm long, yellow, tubular, the tube split to the base, with 5 lobes, 2-lipped. Flowers on stalks 4–11 cm long, single. Flowering: mainly October–February. Seeds elliptic, 3.5 mm long, brown, wing about 0.2 mm wide, brownish.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Goodenia~elongata (accessed 5 April 2021)

Author: Betty Wood.

 $This identification \ key \ and \ fact \ sheets \ are \ available \ as \ a \ free \ mobile \ application:$





