

Plants of South Eastern New South Wales



Flowering plant. Photographer Richard Hartland, near Genoa, Vic



Flowering plant.

Photographer Don Wood, Booderee National Park, Jervis Bay Territory



Line drawing. b. plant. E Mayfield, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Branched Goodenia

Family

Goodeniaceae

Where found

Shrubland, wet **heath**, damp places, and roadsides. Coast, ranges, and tablelands.

Notes

Annual or **perennial** herb to 0.5 m high. Stems hairy or hairless. Leaves mostly basal, 1.4–10 cm long, 6–10 mm wide, thin, tips pointed, surfaces hairy or becoming hairless, margins toothed. Flowers yellow with purplish centres, 10–14 mm long, tubular, the tube split to the base, with 5 lobes, 2-lipped. Flowers on stalks to 80 mm long, in open clusters on almost leafless stalks. Flowering: chiefly October–April. Seeds brown.

PlantNET description: <http://plantnet.rbg Syd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Goodenia~paniculata> (accessed 6 April 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Creative Commons Attribution 3.0 Australia (CC BY)



Android edition



iOS edition