
Plants of South Eastern New South Wales



Flowers. Australian Flant Image Index, Australian Tropical Rainforest key, photographer B Jago, place unknown



Flower spikes. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra



Fruit. Australian Flant Image Index, photographer Murray Fagg, near Robertson



Flow ering plant. Photographer Don Wood, Australian National Botanic Gardens, Canberra

Common name

Settlers' Twine, Boorgay

Family

Araceae

Where found

Wet forest and wet gullies. Coast and ranges north from west of Ulladulla.

Notes

Perennial herb or shrub to about 2 m tall. Fleshy 'petals' and fruit. Leaves alternating up the stems, usually 100–150 cm long, mostly 10–25 mm wide, tough-fibrous, hairless, strongly veined, channelled, tapering to a point. Individual flowers small, with 4 fleshy 'petals'. Pollen bright yellow. Main axis of the inflorescence 1.5-3 m high. Individual flower spikes to about 120 mm long, at first with the flowers green and clustered, the flowers later darkening and becoming separated from each other. Flowers emerge over a long period of time. Fruit blue to blue-black. Flowering: throughout the year.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Gymnostachys~anceps (accessed 19 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





