# **Plants of South Eastern New South Wales**



Flower. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Young seed cases and leaves. Australian Plant Image Index, photographer Murray Fagg, Woko National Park NW of Gloucester



Seeds in open seed case. Australian Flant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT.



Flowering tree. Australian Flant Image Index, photographer Murray Fagg, Cranbourne Botanic Gardens, Melbourne, Vic

#### Common name

Native frangipani

### **Family**

Pittosporaceae

## Where found

Forest, regrowth, and near streams. Natural occurrences north from the Blue Mountains. Frequently planted and naturalised outside its natural range.

#### **Notes**

Shrub or small tree to 22 m high. Occasionally partly deciduous. Bark greyish, smooth, with short horizontal lines of lenticels. Young stems hairy with simple and glandular hairs, becoming hairless, marked by lenticels and scars. Leaves alternating up the stems, sometimes clustered and more or less whorled, 7–16 cm long, 30–45 mm wide, soft, hairless, sometimes hairy on lower surface, especially along the midrib, margins entire, rarely toothed or lobed in juvenile plants, tips gradually tapering to a point and often with a small abrupt point. Flowers fragrant, at first creamy white, turning yellow with a deep pink or reddish throat with age, tubular, the tube 20-30 mm long, with 5 petals each 20–40 mm long. Flowers in loosely branched clusters 15-20 cm diameter. Flowering: spring. Seeds reddish brown, with a single wing all round.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Hymenosporum~flavum (accessed 19 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





