Plants of South Eastern New South Wales



Flower head. Australian Flant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Back of flower head. Photographer Zoya Akulova, California, USA



Flow ering plant. Photographer Don Wood, near Panbula Lake



Rosette. Photographer Keir Morse, Oregon, USA

Common name

Flatweed, Catsear

Family

Asteraceae

Where found

Dry forest, woodland, grassy areas, disturbed sites and roadsides, and wetland margins. Widespread.

Notes

Introduced perennial herb to 0.8 m tall. Stems often grooved, usually hairless, sometimes bristly. Leaves in a rosette at the base of the plant, 2–30 cm long, 10-40 mm wide, bristly, rarely hairless, margins dissected to toothed or with slightly wavy margins, tips blunt. Flower heads below the 'petals' broad cylindrical, 5-15 mm in diameter, with many yellow 'petals', each 8-16 mm long. Flower heads single or in open clusters of a few flowers. Flowers most of the year.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Hypochaeris~radicata (accessed 19 January, 2021)

Author: Betty Wood.

 $This identification \ key \ and \ fact \ sheets \ are \ available \ as \ a \ free \ mobile \ application:$





