Plants of South Eastern New South Wales




Flow ering branches. Photographer Don Wood, Booderee National Park, Jervis Bay



Fruiting branches. Photographer Don Wood, Monga State Forest east of Braidwood



Shrub. Australian Plant Image Index, photographer M Parkes, unknown place



Line drawings. © 2021 Royal Botanic Gardens Board, Melbourne, Vic

Common name

Native Currant, Sour Currant Bush

Family

Santalaceae

Where found

Forest, heath, and occasionally on rocky sites. Coast, ranges, and tablelands.

Notes

Shrub to 3 m high. Fruit fleshy. Branches almost cylindrical to angular, longitudinally ridged, covered with minute warts. Leaves narrow to triangular, 0.1–0.2 cm long, lost early. Flowers red to brown, yellowish brown, or greenish, with 4-5 hooded 'petals', each 0.6–0.7 mm long. Flowers in elongated clusters of 12–30 flowers. Flowers Summer-Autumn. Fruit greenish, yellowish-green, or greenish-brown, to purple, 6–8 mm long, oval to round, acidic, edible.

Rare Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Leptomeria~acida (accessed 22 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





