
Plants of South Eastern New South Wales



Seed cases. Photographer Russell Best, Murray Sunset National Park, Vic



Hant. Photographer Russell Best, Murray Sunset National Park, Vic



Line drawing. q. seed case. MMoir, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Black Cotton Bush

Family

Chenopodiaceae

Where found

Often a colonizer of disturbed ground. Woodland, shrubland, grassland, disturbed sites, and floodplains. Western Slopes.

Notes

Shrub or perennial herb to 0.50 m tall. Stems sparsely to moderately hairy. Leaves fleshy, alternating up the stems, 0.5-1.1 cm long, narrow cylindrical, hairless. Flowers small, tubular, with 5 'petals', hairless, or the margins of the 'petals' fringed. Flowers single, but crowded into leafy spikes at the ends of the branchlets. Seed cases pink to white or tinged orange, drying dark brown to black, 6–10 mm in diameter, hairless, tubular, the tube shallowly hemispherical, 2.5–3 mm in diameter, with a single horizontal wing with one radial slit

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Maireana~decalvans (accessed 24 January, 2021)

Author: Betty Wood.

 $This \, identification \, key \, and \, fact \, sheets \, are \, available \, as \, a \, \, free \, mobile \, application: \, \\$





