


Plants of South Eastern New South Wales



Flowers and leaves. Photographer Jackie Mies



Flowers and leaves. Photographer Don Wood, Budawang National Park west of Batemans Bay



Leaves. Photographer Jackie Mies

Common name

Hairy Milk Vine, Yellow Milk Vine

Family

Apocynaceae

Where found

Forest and wet gullies. Coast, ranges, and the eastern edge of the tablelands.

Notes

Thin-stemmed woody twiner to 5 m high. Stems hairy with yellowish hairs. Leaves opposite each other, 3–12 cm long, 15–45 mm wide, thick, margins curved down, upper surface dark green, more or less hairless, lower surface pale yellowish-green, densely hairy, **glands** 1–3 at the base of the **midvein**, finger-like, tips abruptly pointed or blunt. Flowers cream to pale yellow, 3–4 mm in diameter, tubular, with 5 lobes, in open clusters. Flowers mainly Summer.

Family was Asclepiadaceae

Rare Vic.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Marsdenia-flavescens> (accessed 24 January, 2021)

Author: Betty Wood.



Creative Commons Attribution 3.0 Australia (CC BY)

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition