


Plants of South Eastern New South Wales



Flowering and fruiting stem. Photographer Don Wood, Eurobodalla National Park



Flowering and fruiting stem. Photographer Jackie Miles



Fruiting stem and leaves. Photographer Jackie Miles

Common name

Boobialla

Family

Scrophulariaceae

Where found

Forest, woodland, heath, coastal cliffs, and sand dunes. Coastal areas.

Notes

Shrub to 2 m high, or sprawling to **prostrate**. Fruit fleshy. Branches hairless, not warty, rarely obscurely warty. Leaves **alternating** up the stems, 1.2–7 cm long, 5.5–36 mm wide, tips pointed or blunt, margins **entire**, surfaces the same colour on both sides, shiny, hairless. Flowers white, unspotted, 6–8 mm in diameter, tubular, with 5 lobes. **Sepals** lacking obvious translucent membranous margins. Flowers in clusters of 1–8. Flowering: spring–summer. Fruit light purple to purple or violet, smooth.

Family was Myoporaceae.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Myoporum-boninense> (accessed 24 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Creative Commons Attribution 3.0 Australia (CC BY)



Android edition



iOS edition