Plants of South Eastern New South Wales



Flowering branches. Australian Flant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Flow ering branches. Photographer Don Wood, Bombala State Forest near Bombala



Flow ering branches. Australian Flant Image Index, photographer Murray Fagg, near Numeralla



Flow ering shrub. Australian Plant Image Index, photographer Murray Fagg, near Numeralla

Common name

Violet daisy-bush

Family

Asteraceae

Where found

Dry forest and woodland, often on rocky outcrops. Kosciuszko National Park. Tablelands and ranges south from west of Moruya. Tinderry Range east of the ACT.

Notes

Shrub to 1.5 m high. Branchlets densely hairy with a mat of whitish hairs, often slightly sticky. Leaves alternating up the stems, scattered or crowded, 0.2–1.2 cm long, 1.5–3.5 mm wide, tips broad-pointed without a mucro; margins entire or rough, rolled down, upper surface hairless, sticky and green, often slightly rough, lower surface whitish- or grey-woolly. Flower heads with 7–15 'petals', white or mauve to violet, and cream, yellow, blue, or purplish centres, 13–30 mm in diameter overall. Flower heads approximately conical below the 'petals'. Flower heads single or a few together, at the ends of the branchlets. Flowering: September–November.

Rare Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Olearia~iodochroa (accessed 23 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





