Plants of South Eastern New South Wales



Plants. Australian Plant Image Index, photographer Colin Totterdell, Kosciuszko National Park



Line drawings. c. plant; 'seed'. K Meling, University of Sydney, © 2021 Royal Botanic Gardens Board, Melbourne, Vic

Common name

Branched Caraway, Rock Carraway, Caraway

Family

Apiaceae

Where found

Alpine herbfields and very low shrubland in alpine feldmark, mainly in rocky areas. Kosciuszko National Park.

Notes

Perennial herb to 0.40 m high. Leaf stalks hairy. Base of the plant covered with the remains of old leaf sheaths. Leaves compound, woolly; basal sheaths woolly, tips pointed. Basal leaves 1.5–8 cm long, 7–40 mm wide, with 7–17 dissected leaflets each 0.3–2.5 cm long. Stem leaves alternating up the stems, smaller. Flowers white or pink with a darker midvein, to deep magenta, with 5 petals each about 1-1.2 mm long, hairy on the outside, in 10–15-flowered clusters. 'Seeds' dark red-brown to black. Flowering: summer.

Vulnerable Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Oreomyrrhis~brevipes (accessed 29 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:







Creative Commons Attribution 3.0 Australia (CC BY)