


# Plants of South Eastern New South Wales



Plants. Australian Plant Image Index, photographer Murray Fagg, unknown place



Plants. Australian Plant Image Index, photographer Colin Totterdell, Kosciuszko National Park



Line drawings. e. plant; 'seed'. K Maling, University of Sydney, © 2021 Royal Botanic Gardens Board, Melbourne, Vic

## Common name

Bog caraway

## Family

Apiaceae

## Where found

**Woodland**, **herbfields**, and in and around bogs. Kosciuszko National Park and ACT. Tablelands and the western edge of the ranges east and south from the ACT.

## Notes

**Perennial** herb to 0.45 m tall, highly aromatic with a sickly sweet odour. Flower stalks hairy. Leaves in a basal **rosette**, **compound** or deeply **dissected**, 1–9 cm long, 5–30 mm wide, with 9 to about 21 leaflets each 0.3–1.5 cm long, 2–10 mm wide, hairless except for the finely toothed to fringed margins, basal **sheaths** hairless, tips pointed and **mucronate**. Flowers white with a yellow **midvein**, with 5 **petals** about 1 mm long, hairless or hairy on the outside, in 5–25-flowered clusters. Flowering: summer. 'Seeds' dark brown.

All native plants on unleased land in the ACT are protected.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Oreomyrrhis~ciliata> (accessed 29 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Creative Commons Attribution 3.0 Australia (CC BY)



Android edition



iOS edition