
Plants of South Eastern New South Wales



Rower (subsp. *setigerum*). Photographer Jackie Mles



Flowering plant. Photographer Don Wood, Princes Highway north of Bega



Line drawing. e. flowering stem, f. seed case. M Moir, © 2021 Royal Botanic Gardens Board, Melbourne, Vic

Common name

Opium poppy, Common poppy

Family

Papaveraceae

Where found

Grassy areas, roadsides, disturbed sites, dunes, beaches and stream banks. Western Slopes. ACT. Coast and ranges north of Berry. Widespread elsewhere but not common.

Notes

Introduced annual herb to 1.5 m tall. Stems hairless or bristly, often glaucous. Leaves basal and alternating up the stems, to 30 cm long, glaucous to sparsely bristly, margins often lobed, usually toothed, bases often cordate, upper leaves stem-clasping. Flowers with 4 white, pink, mauve, red, or purple petals, often with a purple spot at the base, each 40-80 mm long. Flowers Oct.—Apr.

subsp. somniferum: Stems and leaves with no or few bristly hairs. Seed cases round, frequently to 5 cm in diameter, and do not release their seed.

subsp. *setigerum*: Stems and leaves covered in bristly hairs. Seed cases cylindrical to almost round, usually less than 2 cm in diameter, and release the seed by pores.

Subspecies are not recognised in the ACT and Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Papaver~somniferum (accessed 29 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)