Plants of South Eastern New South Wales




Flowers. Australian Flant Image Index, photographer Murray Fagg, Mt Majura Nature Reserve, Canberra, ACT



Flower heads with mostly spent flowers. Photographer Keir Morse, Oregon, USA



Flowering stems. Photographer Don Wood, Tathra



Line drawings. (as *Petrorhagia velutina*) BMMayfield, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Velvet pink

Family

Caryophyllaceae

Where found

Weed of grassy areas and disturbed ground. Widespread.

Notes

Introduced annual herb to 0.6 m high. Stems usually deeply furrowed, the middle part usually densely hairy with glandular hairs. Leaves opposite each other, 1–6 cm long, 1–3.5 mm wide, linear, margins finely toothed, tips pointed, the stem-encircling sheath at the base at least twice as long as broad. Flowers with 5 more or less notched pink to purplish petals 10–13 mm long. Flowers appearing single, in tight heads surrounded by brown bracts, one flower opening at a time. Flowering: September–December.

Sometimes hybridises with Petrorhagia nanteuilii.

Was Petrorhagia velutina.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Petrorhagia~dubia (accessed 31 January, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





