Plants of South Eastern New South Wales



Flowering branch. Photographer Jackie Miles, Jerrys Rains, west of Singleton



Flow ering branch. Photographer Jackie Mles, Jerrys Rains, west of Singleton

Common name

A Rice Flower

Family

Thymelaeaceae

Where found

Coast, ranges, and eastern tablelands, north from west of Nowra. Near Jindabyne. Possibly ACT.

subsp. elliptifolia: Rocky sites. Western Sydney to west of the Blue Mountains. Mostly north of the area covered by this key.

subsp. hirsuta: Forest. Coast, ranges, and eastern tablelands, north from west of Nowra. Near Jindabyne. Possibly ACT.

Notes

Shrub to 0.5 m high. Tough 'bootlace' bark when the stems broken. Stems hairy. Leaves opposite each other, sometimes alternating up the stems, 0.2–3.5 cm long, 1–11 mm wide, hairy at least on the lower surface, medium to dark green or blue-green on the upper surface, paler below. Flowers bisexual or female, greenish yellow, or yellow to white, 6–12 mm long, tubular, with 4 spreading lobes each 1–3.5 mm long, often red, rarely white. Flowers in 1–18 flowered heads. Bracts absent or similar to the leaves. Flowers Apr.–Oct.

subsp. *elliptifolia*: Leaves 0.2–3.5 cm long, 2–11 mm wide, covered on both surfaces with fine or coarse long hairs, margins fringed. Flowers 3.5–8 mm long, usually yellow or greenish yellow and sometimes with red lobes, rarely white. Heads 5–18-flowered.

subsp. *hirsuta*: Leaves covered on both surfaces with short fine, slightly tangled, hairs. Leaves 0.2–1.5 cm long, 1–6 mm wide, margins fringed. Flowers 4–7 mm long, yellow with dark red lobes. Heads 1–7-flowered.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Pimelea~latifolia (accessed 31 January, 2021)

Author: Betty Wood.

 $This identification \ key \ and \ fact \ sheets \ are \ available \ as \ a \ free \ mobile \ application:$







Creative Commons Attribution 3.0 Australia (CC BY)