Plants of South Eastern New South Wales



Flow ering stem Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra



Flow ering branches. Photographer Neville Walsh, © 2021 Royal Botanic Gardens Board, Melbourne, Vic



Line drawing. f. leaf. A. Barley, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Blunt-leaf pomaderris

Family

Rhamnaceae

Where found

Rocky places and near streams. West of Batlow. Tablelands and western edge of the ranges between the Hume Highway and south of Queanbeyan. In Victoria along the NSW border.

Notes

Shrub to 2 m high. Stems greyish to rusty stellate hairy (needs a hand lens or a macro app on your phone/tablet to see), rarely with scattered simple hairs. Leaves alternating up the stems, 0.5–4.5 cm long, 1–10 mm wide, upper surface bristly with scattered simple hairs; lower surface greyish and densely stellate-hairy, veins covered with long simple hairs, lateral veins obscure or looping to the inside and not reaching the leaf margins, tips blunt, margins curved down. Flowers cream or yellow, with 5 sepals each 1.5–2 mm long, falling early, and 0 petals, in branched clusters, usually about 20–50-flowered.

Rare Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Pomaderris~helianthemifolia (accessed 2 February, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





