Plants of South Eastern New South Wales



Flowers, buds, and leaf. Australian Hant Image Index, photographer Murray Fagg, near Binalong, NW of Yass



Flowers. Australian Plant Image Index, photographer Murray Fagg, near Binalong, NW of Yass



Line drawing. j. flower cluster; k. leaf. BMMayfield, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Sulphur cinquefoil

Family

Rosaceae

Where found

Pastures, roadsides, and weedy areas. Tablelands, ACT, the mountains to the west, Kosciuszko National Park, and Western Slopes.

Notes

Introduced perennial herb to 0.7 m high. Stems green or reddish, sparsely covered with short and long hairs. Flowering stems densely hairy. Leaves basal and alternating up the stems, 5-12 cm long, compound, with 5-7 leaflets each 1.5-8 cm long, 5-20 mm wide, upper surface almost hairless, lower surface sparsely hairy with white spreading hairs, margins toothed. Flowers yellow, with 5 petals each 6–12 mm long, in many-flowered open clusters. Flowers Spring to summer.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Potentilla~recta (accessed 3 February, 2021)

Author: Betty Wood.



This identification key and fact sheets are available as a free mobile application:





Creative Commons Attribution 3.0 Australia (CC BY)