

Plants of South Eastern New South Wales



Cluster of flower heads. Photographer Richard Hartland, Paddys Ranges near Maryborough, Vic



Flowering stems. Photographer Don Wood, East Boyd State Forest south of Eden



Plants. Photographer Richard Hartland, Paddys Ranges near Maryborough, Vic



Painting from Jacob Sturm (1796) *Deutschlands Flora in Abbildungen* Wikimedia

Common name

Jersey cudweed

Family

Asteraceae

Where found

Variety of **habitats** including **forest**, **woodland**, and open disturbed sites, often near water. Widespread.

Notes

Annual herb to 0.55 m tall. Stems **woolly**. Basal leaves withering early. Stem leaves **alternating** up the stems, 1–6.5 cm long, 1–8 mm wide, bases slightly stem claspings, surfaces hairy, margins sometimes curved to rolled down, often wavy. Flower **heads** about 3–4 mm long, 4–6 mm in diameter, urn-shaped to oval. **Florets** cream to yellow, with 0 **petals**, in clusters of about 5–20. Flowers most of year.

Was *Helichrysum luteoalbum*.

Laphangium luteoalbum in Vic.

All native plants on unleased land in the ACT are protected.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Pseudognaphalium~luteoalbum> (accessed 1 February, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)