Plants of South Eastern New South Wales




Flowering stem. Photographer Arthur Chapman, near Ballarat, Vic



Flowering stems. Photographer Jackie Miles, Burrowa-Pine Mountain National Park, Vic.



Shrub. Photographer Arthur Chapman, near Ballarat, Vic



Line drawings. c. flowering stem leaves and stipules; calyx. J Armstrong, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Flat-leaf Bush-pea

Family

Fabaceae

Where found

Dry forest. Mainly east of Albury. Rarely elsewhere.

Notes

Shrub to 2 m high. Stems sparsely to densely appressed-hairy. Leaves alternating up the stems, 0.8–2 cm long, 3–6 mm wide, concave in cross section, tips rounded and usually minutely notched, margins flat, upper surface becoming hairless, darker than the lower surface, lower surface hairless. Flowers 7–12 mm long, pea shaped, with 5 petals, 2 joined together to form the keel. Standard petal yellow to orange, wings yellow to orange, keel red to purple. Bracteoles linear, hairy, inserted on the calyx tube. Flowers in dense clusters of more than 5 flowers. Flowers September-November. Pods densely to sparsely hairy, smooth.

Rare Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Pultenaea~platyphylla (accessed 2 February, 2021)

Description partly taken from: Kok, R.P.J. de & West, J.G., (2004) A revision of the genus *Pultenaea* (Fabaceae) 3. The eastern species with curved down leaves. *Australian Systematic Botany* 17(3): 307-308, Fig. 14 (map)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)