# **Plants of South Eastern New South Wales**




Flow ering stems. Photographer Don Wood, east of Braidwood



Flow ering stems. Photographer Don Wood, Wadbilliga National Park



Shrub. Australian Plant Image Index, photographer M Orisp, Mongarlowe, east of Braidwood



Line drawings. h. flowering stem leaves and stipules; flower. J Armstrong, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

## Common name

Low bush-pea

### Family

Fabaceae

#### Where found

Wet to dry forest, woodland, heath, and disturbed sites, usually in stony areas. Widespread. Rarely coastal.

#### **Notes**

Shrub, prostrate or to 0.6 m tall. Stems hairy to sparsely hairy. Leaves alternating up the stems, 0.2-1 cm long, 1-2 mm wide, concave in cross section, tips blunt, margins curved to rolled upwards, upper surface hairless, paler than the lower surface or both surfaces the same colour; lower surface covered in minute warts, with scattered, pale hairs. Flowers pea shaped, with 5 petals, 2 joined together to form the keel. Standard petal 7.6-9.5 mm long, yellow to pink to orange-red, sometimes with red to red-brown markings. Wings yellow or pink to orange-red. Keel orange to red-brown. Bracteoles with 3 teeth at the tips, papery, inserted at the base of the calyx tube. Flowers in leafy clusters. Flowers most of the year. Pods with a tuft of hairs at the tips.

All native plants on unleased land in the ACT are protected.

Vulnerable Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Pultenaea~subspicata (accessed 4 April 2021)

Description partly taken from: Kok, R.P.J. de & West, J.G., (2002) A revision of *Pultenaea* (Fabaceae) 1. Species with ovaries hairless and/or with tufted hairs. *Australian Systematic Botany* 15(1): 108-109

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)