
Plants of South Eastern New South Wales



Flow ering stems. Australian Flant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Flow ers and leaves. Photographer Richard Hartland, Lake Mountain, Vic



Line drawings. g. leaf; petal; petal base and nectary; seed. BMMayfield, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Strawberry Buttercup

Family

Ranunculaceae

Where found

Woodland, grassland, and moist areas, and bogs. Kosciuszko National Park, ACT, and Blue Mountains.

Notes

Perennial herb, stoloniferous. Flower stalks to 0.22 m high, hairy to almost hairless, with appressed or spreading hairs. Leaves basal, 0.3–4 cm long and 3–50 mm wide, round or almost round, usually compound, sometimes deeply lobed, divided into 3 leaflets or lobes which are toothed or further lobed, surfaces dark-green, more or less shining above, paler beneath, both surfaces hairless to moderately appressed-hairy. Flowers bright golden-yellow, glossy, with 5–11 petals each 5-10 mm long, Sepals spreading, rarely turned back. Flowering stems 1–2 flowered. Flowering: spring to summer. Seeds 8–30, 2–2.5 mm long, broadly oval, with 2–4 oblique coarse low ridges, or with a bold network on each face; beak 0.5–1 mm long, curved down.

Hybrids between Ranunculus millanii and Ranunculus collinus have been given the name Ranunculus x ligulatus.

All native plants on unleased land in the ACT are protected.

Rare Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Ranunculus~collinus (accessed 4 February, 2021) Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)