
Plants of South Eastern New South Wales



Flower. Photographer Keir Morse, California, USA



Flow ering plants. Photographer Jan Sevcik, Czech Republic



Line drawings (as Batrachium trichophyllum). flowering plant; petal; petal base and nectary; seed BMMayfield, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

A buttercup

Family

Ranunculaceae

Where found

Notes

Introduced submerged perennial herb, only the flowers emerging above the water. Stems to about 0.6 m long, often rooting at the lower nodes. Leaves alternating up the stems, 1–5 cm long, 10-50 mm wide, divided into numerous thread-like segments 0.1–0.5 mm in diameter, not lying in the same plane, segments often collapsing together when withdrawn from the water, bases dilated into membranous, often appressed-hairy sheaths. Flowers entirely white, or yellow at the base, with 4-5 petals each 3–7 mm long. Sepals greenish, often tinged purple, curved back or spreading, soon falling. Flowers single. Flowers Oct.—Feb. Seeds 15–35, lens-shaped to oval, usually sparsely hairy, 1.5-2 mm long, sides prominently transversely ridged; beak absent or to 0.2 mm long.

Was Batrachium trichophyllum.

Treated as Introduced in NSW. Uncertain Status (Native or Introduced) in Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Ranunculus~trichophyllus (accessed 16 April 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





