Plants of South Eastern New South Wales



Flowering stems. Australian Flant Image Index, photographer Murray Fagg, near Bredbo



Flow ering plant. Photographer Don Wood, Namadgi National Park, ACT



Flow ering plant. Photographer Don Wood, Booderee National Park, Jervis Bay Territory

Common name

Erect Groundsel

Family

Asteraceae

Where found

Forest, woodland, rocky areas, near swamps and rivers, and other moist sites. Widespread.

Notes

Annual or perennial herb to 1.5 m high. Stems hairless or sparsely rough hairy near the base. Leaves alternating up the stems, 5-18 cm long, 4–50 mm wide, hairless or nearly so, margins usually with 3–8 triangular lobes per side, irregularly dissected, toothed, or entire, bases with small to moderate auricles on the leaves above the mid-stem. Flower heads cylindrical, 4.5-6.5 mm long, with 0 petals, surrounded by 7–11 appressed green bracts. Florets 8–25, about 2mm longer than the flower head. Flower heads often in 50-200 flowered clusters. Flowering: mainly summer–autumn.

A feature of this species is the precociously developed leafy growth at the bases of the leaves.

All native plants on unleased land in the ACT are protected.

Rare Vic.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Senecio~diaschides (accessed 6 February, 2021) Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)