


Plants of South Eastern New South Wales



Flowering plant. Australian Plant Image Index, photographer AN Schmidt-Lebuhn, Illunie Nature Reserve south of Cowra



Flowering plants. Australian Plant Image Index, photographer Murray Fagg, Scottsdale Bush Heritage Reserve near Bredbo



Line drawings. a. leaf; seed. Espeth Beckett, University of Oxford., © 2021 Royal Botanic Gardens Board, Melbourne, Vic

Common name

Jo-jo, Bindyi, Lawn Burweed

Family

Asteraceae

Where found

Weed of lawns and playing fields, disturbed sites, and rocky areas. Widespread, mainly in the Sydney area and Canberra.

Notes

Introduced annual herb, **prostrate**, rooting at the **nodes**. Burrs with many seeds, each seed with thin, broad-lobed **wings**, hairless or almost hairless at the top, with a spine 1.5-3 mm long. Stems branched, **spreading** or absent, sparsely hairy. Leaves basal and in clusters, 1.5–5 cm long, 5–15 mm wide, carrotlike, **dissected**, hairy, bases sheathing. Flower **heads** with 0 **petals**, 4–10 mm in diameter at the seeding stage, slightly convex, single at the base of each leaf cluster. Flowering: spring–summer.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Soliva-sessilis> (accessed 8 February, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)